

(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 108 425 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
08.06.2005 Bulletin 2005/23

(51) Int Cl.7: A61K 9/50, A61K 31/44

(21) Application number: 99670010.0

(22) Date of filing: 16.12.1999

(54) **New stable multi-unitary pharmaceutical preparations containing substituted benzimidazoles**

Neue stabile mehrereinheitliche substituierte Benzimidazole enthaltende pharmazeutische Präparate

Nouvelles préparations pharmaceutiques multi-unitaires stables contenant des benzimidazoles substitués

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(43) Date of publication of application:
20.06.2001 Bulletin 2001/25

(73) Proprietor: Laboratório Medinfar-Produtos
Farmaceuticos, S.A.
2700 Amadora (PT)

(72) Inventors:
• Gonçalves Mendes, Carla Patricia
2685 Sacavem (PT)
• Caeiro Ramalho De Oliveira, Maria Julia
2685 Sacavem (PT)

(74) Representative: Pereira da Cruz, Joao
J. Pereira da Cruz, S.A.
Rua Vitor Cordon, 14
1249-103 Lisboa (PT)

(56) References cited:

EP-A- 0 277 741	EP-A- 0 519 144
EP-A- 0 773 025	WO-A-92/22284
WO-A-98/19668	US-A- 5 385 739
US-A- 5 817 338	

- DATABASE WPI Section Ch, Week 199744
Derwent Publications Ltd., London, GB; Class
B02, AN 1997-478104 XP002138129 & PT 101 826
A (LAB MEDINFAR PROD FARM SA) , 30
September 1997 (1997-09-30)
- DATABASE WPI Section Ch, Week 199844
Derwent Publications Ltd., London, GB; Class
B02, AN 1998-518792 XP002138130 & PT 101 965
A (LAB MEDINFAR PROD FARM SA) , 30
September 1998 (1998-09-30)

Remarks:

The file contains technical information submitted
after the application was filed and not included in this
specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 108 425 B1

Description

Field of the Invention

[0001] The present invention relates to new oral multi-unitary pharmaceutical preparations containing substituted benzimidazoles being inhibitors of H⁺, K⁺-ATPase (i.e. omeprazole, lansoprazole, pantoprazole, leminoprazole and pariprazole) or their pharmaceutically acceptable salts. The preparations comprise a spherical inert core, constituted by starch and sugar, coated with a layer containing at least one substituted benzimidazole in the micronized form, which is mixed with pharmaceutically acceptable inert excipients, whose proportions are suitable for allowing the disaggregation of the dosage forms and the intended dissolution of the active ingredient(s), this layer in turn being coated with an insulating layer of an exclusively polymeric nature and of suitable thickness, applying lastly an external, gastroresistant or enteric layer of suitable thickness, in order to guarantee the integrity of the product until it reaches the proximal part of the small intestine, where the formulation will be disaggregated to facilitate the absorption of the substituted benzimidazolic compounds. The pellets produced according to the invention are placed in hard gelatine capsules and administered to patients in this form. The invention does not require the stabilization of the benzimidazolic compounds using any of the strategies or processes which already belong to the state of the art. Besides, the pharmaceutical products produced in this way are free from organic solvents and/or the impurities generally associated to them, because the application of the different layers exclusively requires aqueous solvents. This aspect constitutes a technological advantage, since the aspect constitutes a technological advantage, since the manufacturing process is incomparably safer, because there is no toxicity risk to operators or explosion risks and moreover it is more ecological, because there is no possibility of environmental contamination caused by organic solvents leaking into the atmosphere. Finally, it is much safer for the patient, because there is no need to consider solvents and/or residual impurities associated to them, which is a considerable advantage in terms of public health. For the manufacture of the products, only equipment is required - a fluid bed equipment with an inner partition device (wurster). The products obtained by extrusion/spheronization, by roto granulation or by "powder coating" are completely outside the scope of the present invention.

[0002] The products obtained are stable for a period of time compatible with pharmaceutical requirements, and they present gastroresistance $\geq 85\%$ and dissolution characteristics generally adapted to the period of validity established for pharmaceutical products (i.e. 3 years).

Background of the Invention

[0003] Benzimidazolic compounds such as omeprazole (5-methoxy-2-(((4-methoxy-3,5-dimethyl-2-pyridinyl)-methylsulfinyl)1H-benzimidazole (EP-B1-0005129), lansoprazole (2-(((3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridyl)methyl(sulfinyl) 1H-benzimidazole) (US-P-4,628,098), pantoprazole (US-P-4,758,579)), leminoprazole and pariprazole are anti-ulcerous substances known for decreasing gastric acid secretion (Olbe L., *et al.*, *Gastroenterol.*, **83**:193-198 (1982); Saton H. *et al.*, *Jpn. J. Pharmacol.* **40** (suppl.), 226 (1986); Saton H. *et al.*, *J. Pharmacol. Exp. Ther.*, **248** (2), 806-815 (1989), Nagaya, H. *et al.*, *J. Pharmacol. Exp. Ther.*, **248** (2), 799-805 (1989)) and they are used in the therapeutics of diseases related to gastric acidity in mammals and especially in humans, including gastric and duodenal ulcers, reflux oesophagitis, gastritis, duodenitis and Zollinger-Ellison syndrome.

[0004] It is also known that these substituted benzimidazoles possess a very low level of solubility in water, they solubilize easily in alkaline solutions (i.e. the $t_{1/2}$ of degradation of omeprazole in aqueous solutions of pH>11 is 600 days), they degrade quickly in acidic and neutral media (i.e. the $t_{1/2}$ of degradation of omeprazole in aqueous solutions of pH<4 is 10 minutes) (Pilbrand and Cederberg, *Scand. J. Gastroenterol.*, **20** (Suppl. **108**), p. 113-120 (1985)) and they are stabilized in the presence of alkaline reacting compounds. The maximum stability of omeprazole solutions is reached at pH 11; for pHs lower than 7.8, degradation occurs very quickly (*Drug.Dev.Ind.Pharm.*, **21**(8), 965 (1995)). Lansoprazole is relatively stable when exposed to the light. This compound degrades in aqueous solution, its degradation speed increasing as the pH decreases. For this reason, whenever the oral route is used it is indispensable to coat the formulations with a gastroresistant layer, in order to avoid contact between the active ingredient and gastric content with acidic characteristics. On the other hand, it is necessary that the formulations dissolved quickly in the intestine, where the benzimidazolic compounds should be absorbed, i.e. when the pH becomes higher than 6.8. Several pharmaceutical preparations containing substituted benzimidazole inhibitors of H⁺,K⁺-ATPase are already patented.

[0005] The first assays performed with classic enteric pellets containing omeprazole showed that they did not present suitable stability during a long enough period of time to be used in a satisfactory pharmaceutical dosage form, because fast degradation of the active ingredient was observed when subjected to normal conditions of storage (T=25±1°C and HR=40-75%), with the appearance of noxious degradation products (Brandstrom A., *et al.*, *Acta Chemica Scandinavia*, **43**, 536-548 (1989)).

[0006] DE-A1-3046 559 describes a way to coat a pharmaceutical dosage form with an insoluble layer containing

microcrystalline cellulose and with a second enteric layer. This preparation did not allow the desirable release of omeprazole in the small intestine.

[0007] US- A - 2 540 979 describes a preparation coated with an enteric layer combined with a first coating of a water-insoluble wax. This preparation was shown to be inadequate for omeprazole because the direct contact of the cellulose acetophthalate (CAP) with the omeprazole caused its degradation.

[0008] DE-A1-1 204 363 describes a pharmaceutical dosage form which has three layers but cannot be used for omeprazole because it does not allow the fast release of omeprazole in the small intestine, as is desirable. DE-A1 - 1 204 363 concerns a coating that allows the delivery of the active ingredient in the ileum, which is not desirable in the case of omeprazole.

[0009] DE-A1-1 617 615 describes a dosage form which likewise cannot be applied to omeprazole because it is dissolved very slowly in the intestinal juice.

[0010] GB 2 189 698 A and US 4 786 505 propose the mixture of a mass of cellulose derivatives and disaggregants, with an appropriate amount of omeprazole and alkaline salts, in order to obtain, by extrusion, a core which is spheronized and coated with gastroresistant agents dissolved in alcoholic solutions also containing considerable percentages of acetone. However, the pellets obtained are extremely irregular in shape and dimensions, which hinders the process of filling the gelatine capsules, and this can have repercussions on a relative dispersion of the average weight of the capsules and of the respective dosage. Besides, the use of organic solvents in any of the phases of preparation of the pharmaceutical dosage forms means that the quantity thereof in the final product will have to be determined, as well as the quantity of the impurities that may be associated to them and may be potentially dangerous for the health, which makes them highly inadvisable.

[0011] US-5 232 706 describes omeprazole dosage forms coated with one or more layers containing water-soluble alkaline excipients mixed with a second alkaline compound.

[0012] US-5 385 739 and FR-A-2 692 146 describe pharmaceutical gastroprotected dosage forms containing omeprazole wherein the active substance is diluted with mannitol and applied to an inert core. The dosage form is prepared in a conventional coating pan, which is a slow process and requires the use of organic solvents which are noxious to the operator and dangerous to handle. Besides, with the pellets obtained, with a wrinkled surface, it is difficult to fill capsules of uniform weight. Finally, the pellets marketed by the applicant of these patents did not observe the gastroresistance and dissolution criteria required for their use during the period of validity established for pharmaceutical dosage forms.

[0013] US-5 399 700 concerns the stabilization of benzimidazolic derivative compounds as cyclodextrin inclusion complexes.

[0014] US-5 690 960 concerns dosage forms containing magnesium omeprazole salts with more than 70% crystallization.

[0015] US-5 708 017 concerns an oral formulation in the form of a paste in an oily vehicle, containing a proton pump inhibitor, intended for the treatment of gastric ulcers in humans and animals.

[0016] US-5 714 504 relates to pharmaceutical formulations for oral administration containing alkaline salts of the (-) enantiomer of omeprazole.

[0017] WO 96/01623 relates to omeprazole tablets obtained from gastroresistant multi-unitary dosage forms, which are used to prepare tablets in the presence of appropriate excipients.

[0018] WO 97/12581 relates to oral formulations of omeprazole obtained by compression of pellets containing benzimidazoles, thereby obtaining micro-tablets which are subsequently coated and placed in hard gelatine capsules.

[0019] WO 98/53803 relates to the specific use of a quality of HPMC in the manufacture of pharmaceutical dosage forms containing omeprazole.

[0020] WO 98/19668 relates to an omeprazole formulation with pulsatile kinetics which simulates a bolus for a once-a-day dosage form.

[0021] WO 98/50019 relates to oral dosage forms containing omeprazole or lansoprazole, inside hard gelatine capsules which are subsequently coated with gastroresistant or enteric polymers.

[0022] US-4 689 333 describes pharmaceutical compositions for the prevention or treatment of ulcers containing an effective amount of lansoprazole or a pharmaceutically acceptable salt thereof.

[0023] US-5 026 560 relates to spherical pellets consisting of a core coated with a binding agent which is coated with a mixture of powders containing lansoprazole, HPMC of low viscosity and magnesium or calcium carbonates as alkaline agents. The manufacturing process used was "powder coating", which is difficult and laborious and therefore makes the product very expensive.

[0024] US-5 045 321 relates to dosage forms for the production of pellets or coated tablets which comprise a uniform mixture of lansoprazole with at least one alkaline inorganic salt. There is no reference to protective and/or enteric coatings in the claims of the patent.

[0025] US-5 093 132 claims a stable oral dosage form containing lansoprazole or a pharmaceutically acceptable salt thereof mixed uniformly with an alkaline inorganic salt as a stabilizing agent. Only a simple reference is made to

the use of an enteric coating.

[0026] EP 0 589 981 relates to oral dosage forms of pantoprazole as tablets or pellets which use alkaline and/or alkaline-earthly metallic salts for stabilization of the active ingredient and which use organic solvents (i.e. isopropanol) in the technological process for the production of the pharmaceutical formulations.

[0027] WO 97/12580 describes oral dosage forms of substituted benzimidazoles, preferably pantoprazole, obtained by compression of pellets containing benzimidazole, thereby producing micro-tablets that are subsequently coated and placed inside hard gelatine capsules.

[0028] PT 101 826 and PT 101 965 disclose pellet formulations of omeprazole constituted by an inert core, an active layer, an isolation layer with a thickness of at least 4 μm and an enteric layer with a thickness of at least 20 μm .

[0029] The present invention concerns stable multi-unitary dosage forms of gastroresistant pellets for oral administration, free from residues of organic solvents and the impurities associated to them, containing a substituted benzimidazole or a pharmacologically acceptable salt thereof, or associations of substituted benzimidazoles or of their pharmaceutically acceptable salts. The formulations do not comprise any of the methods of stabilization of benzimidazoles already known from the state of the art and they are the first ones whose manufacturing process involves the use of a fluid bed equipment with an inner partition device (wurster), and not extrusion/spheronization techniques, roto granulation, "powder coating" or coating in conventional coating pans. Thus, the present invention describes pellet formulations composed of an inert core coated with a layer which contains the active ingredient(s), coated in turn with an intermediate layer of a minimum thickness of 15 μm and with a final external gastroresistant or enteric layer of a minimum thickness of 30 μm . The thickness of the intermediate layer is decisive to the stability of the final product and it prevents the incorporation of alkaline and/or alkaline-earthly metallic salts into the mixture containing the benzimidazolic compounds, as well as into the insulating layer. The pellets obtained have spherical symmetry and their surface is perfectly flat when observed by scanning electronic microscopy (S.E.M.), contrary to the other formulations on the market that are obtained by other methods. In addition, the fact that the products are manufactured without the use of organic solvents makes them safer for patients and for operators who manufacture and handle them. The manufacturing process does not cause environmental pollution problems, which are becoming increasingly essential to avoid, and this is another of the aspects that differentiate these formulations from those belonging to the state of the art. Moreover, the process for manufacturing these new formulations has technical as well as economic advantages. In fact, all the stages of the manufacturing process use the same equipment, which represents less contact for the operator with the product, contrary to what happens with the other described processes which require the transfer of the pellets from the extruder/spheronizer to the coating pans or fluid bed equipments in order to be coated. Neither does it involve various equipments of restricted use and a high cost, which would have to be reflected in the final price of the product to consumers. These facts, together with the high market prices of benzimidazolic compounds, lead to an exaggerated increase in the price of the final product, which will necessarily be borne by patients who need the medication. The present invention seeks to reduce the production costs of these products in order that they can be used on a large scale all over the world.

Summary of the Invention

[0030] The present invention describes multi-unitary pharmaceutical dosage forms containing a substituted benzimidazole (omeprazole, lansoprazole or pantoprazole, leminoprazole or pariprazole) or a salt thereof, or associations of benzimidazoles or their salts, constituted by a succession of layers arranged around an inert, spherical core, prepared in a fluid bed equipment with an inner partition device (wurster). The component elements of the dosage form are:

- (1) inert core,
- (2) active layer,
- (3) insulating layer and
- (4) gastroresistant or enteric coating layer.

[0031] The inert cores of dimensions of 600 - 710 μm , 710 - 850 μm or 850 - 1000 μm , constituted by pharmaceutically acceptable inert excipients are coated with a layer containing at least one benzimidazole or its salts, mixed with pharmaceutically acceptable inert excipients, so that this layer is quickly disaggregable. This layer is then coated with an insulating layer of a strictly polymeric nature, soluble in water of a minimum thickness of 15 μm , and finally this layer is coated with an aqueous enteric coating layer of a minimum thickness of 30 μm . The pellets have spherical symmetry and a flat surface, are free from residues of organic solvents and the impurities associated to them and they have a moisture level that guarantees good stability under normal storage conditions. The pellets are placed in hard gelatine capsules and it is in this form that they are administered to patients. The formulations with this composition are characterized in that they do not dissolve in an acid medium, but dissolve quickly at an alkaline pH and present good stability in terms of dosage and in gastroresistance and dissolution assays, when stored for at least 3 years.

Brief Description of the Figures

[0032] In order to test the stability of the finished products in their packaged form, industrial batches of omeprazole and lansoprazole pellets were prepared (i.e. 400,000 capsules/batch), which were placed inside hard gelatine capsules, which were in turn stored in polyamide/aluminium/PVC blisters sealed with aluminium foil of 25 μm . The blisters were incubated in climatic chambers at 25°C and 60% relative humidity (atmospheric conditions) and 40°C and 75% relative humidity (accelerated assays). The following parameters were periodically determined: dosage, gastroresistance, dissolution and level of impurities.

[0033] Figure 1 shows a graph of the dosage, gastroresistance and dissolution results obtained for omeprazole pellets stored inside blisters and kept at 40°C and 75% relative humidity (accelerated assays) for 9 months. As can be seen, the omeprazole content is never lower than 95%, which is the minimum amount generally admissible for the amount of active ingredient in pharmaceutical forms during the period of validity. As can also be seen in Figure 1, the gastroresistance value obtained for the period during which the assay was carried out was never lower than 85%, which is the minimum percentage generally accepted for gastroresistant formulations during the period of validity. Also as regards the dissolution assay, the value determined for this parameter for the period during which the assay was carried out was never lower than 75%, which is the minimum value generally accepted for this type of formulation during the period of validity. The values obtained for each of the parameters studied are presented in detail in Table V.

[0034] Figure 2 shows a graph of the dosage, gastroresistance and dissolution results obtained for omeprazole pellets stored inside blisters and kept at 25°C and 60% relative humidity (assays in real time or under atmospheric conditions) for 18 months. As can be seen, the omeprazole content is never lower than 95%, which is the minimum amount generally admissible for the dosage of active ingredient in pharmaceutical forms during the period of validity. As can also be seen in Figure 2, the gastroresistance value obtained for the period during which the assay was carried out was never lower than 85%, which is the minimum percentage generally accepted for gastroresistant formulations during the period of validity. Also as regards the dissolution assay, the value determined for this parameter for the period during which the assay was carried out was never lower than 75%, which is the minimum value generally accepted for this type of formulation during the period of validity. The values obtained for each of the parameters studied are presented in detail in Table VI.

[0035] Figure 3 shows a graph of the dosage, gastroresistance and dissolution results obtained for lansoprazole pellets stored inside blisters and kept at 40°C and 75% relative humidity (accelerated assays) for 6 months. As can be seen, the lansoprazole content is never lower than 95%, which is the minimum amount generally admissible for the amount of active ingredient in pharmaceutical forms during the period of validity. As can also be seen in Figure 3, the gastroresistance value obtained for the period during which the assay was carried out was never lower than 85%, which is the minimum percentage generally accepted for gastroresistant formulations during the period of validity. Also as regards the dissolution assay, the value determined for this parameter for the period during which the assay was carried out was never lower than 75%, which is the minimum value generally accepted for this type of formulation during the period of validity. The values obtained for each of the parameters studied are presented in detail in Table VII.

[0036] Figure 4 shows a graph of the dosage, gastroresistance and dissolution results obtained for lansoprazole pellets stored inside blisters and kept at 25°C and 60% relative humidity (assays in real time or under atmospheric conditions) for 12 months. As can be seen, the lansoprazole content is never lower than 95%, which is the minimum amount generally admissible for the dosage of active ingredient in pharmaceutical forms during the period of validity. As can also be seen in Figure 4, the gastroresistance value obtained for the period during which the assay was carried out was never lower than 85%, which is the minimum percentage generally accepted for gastroresistant formulations during the period of validity. Also as regards the dissolution assay, the value determined for this parameter for the period during which the assay was carried out was never lower than 75%, which is the minimum value generally accepted for this type of formulation during the period of validity. The values obtained for each of the parameters studied are presented in detail in Table VIII.

Detailed Description of the Invention

[0037] The present invention describes formulations that comprise an inert core of spherical symmetry constituted by pharmaceutically acceptable inert excipients, coated with a succession of layers, in order to produce final multi-unitary gastroresistant formulations (pellets) that are placed in hard gelatine capsules. Another advantage of these new formulations is that all the stages of their production always take place in the same equipment and do not require the use of organic solvents. For this reason, these pharmaceutical preparations are safer for operators during their manufacture and for the patients to whom they are administered.

[0038] The elements to consider in the formulations are the following:

- I. Inert cores.

- II. Active layer.
- III. Insulating layer.
- IV. Gastroresistant or enteric layer.

I. Inert cores

[0039] The inert cores have dimensions of 600 - 710 μm , 710 - 850 μm , or 850 - 1000 μm , according to the benzimidazole potency that is intended in the final product. The inert cores are constituted by starch (10-30%) and sucrose (60-90%), but can also contain glucose, lactose, mannitol or any other pharmaceutically acceptable inert excipients. With solid consistency and spherical symmetry, the inert cores are obtained from sucrose particles by coating them with successive layers of sucrose and cornstarch. The final product must contain less than 4% humidity. The particles must be selected in a relatively narrow range of granulometry, so that the pellets obtained at the end are of a uniform dimension. For the benzimidazole dosages generally used in therapeutics (i.e. generally 20-40 mg/capsule), the inert cores should not exceed 850 μm , at maximum 1000 μm ; so that gelatine capsules of a relatively small dimension can be used, which helps the patient to adhere to the therapy being followed. The inert cores can be obtained commercially and do not constitute the aim of the present invention.

II. Active layer

[0040] The active layer is applied to the inert core and it contains at least one micronized benzimidazole (i.e. omeprazole, lansoprazole, pantoprazole, leminoprazole or pariprazole) or its pharmacologically acceptable salts and excipients that are also pharmaceutically acceptable, for example surfactant agents, such as, sodium triaurylphosphate, sodium dodecyl sulfate, sodium tridecylethoxylate, sodium myristylsulfate, or, preferentially, fatty acids of sorbitan esters, such as glycerol polyethyleneglycol oxystearate (Cremophor® of BASF Corporation), which increase the wettability of the benzimidazolic compounds; binding agents of a polymeric nature, such as hydroxymethylpropylcellulose (Pharmacoat® of Shinetsu; Methocel® of Colorcon), hydroxypropylcellulose (Klucel® of Hercules), hydroxymethylcellulose, hydroxyethylcellulose (Natrosol® of Hercules or Cellosize® of Union Carbide Corporation), methyl cellulose (Methocel® A of Colorcon), ethylcellulose (Aquacoat® de FMC or Surelease® of Colorcon), polyvinylpyrrolidone (Povidone® of GAF Corp., Kollidon® of BASF), polyethylene glycols, etc. The mixture that constitutes the layer also contains sodium tetraborate to improve the solubility and stability of the benzimidazoles. In order for the dosage form to dissolve quickly disintegrating agents are added, such as sodium starch glycolate (Explotab® of Mendell, Primogel® of Generichem), pregelatinized starch (Starch 1500), microcrystalline cellulose (Avicel of FMC), sodic or calcic carboxymethylcellulose or sodium alginate. Lubricant agents such as magnesium or calcium stearate, or preferentially talc, should be added, in order to decrease the friction between the particles. The suspension containing the mixture of the ingredients is prepared exclusively in water. For the suitable preparation and application of the suspension, the inclusion of anti-foaming agents such as simethicone is advisable, preferentially in a form which is easily dispersible in water (in the form of an emulsion containing not less than 85% polydimethylsiloxane).

[0041] When used in the appropriate proportions, the substances previously mentioned are mixed with the active ingredient in an aqueous solution, are atomized on the inert cores under suitable conditions, producing a uniform layer of a thickness of between 150 and 250 μm .

III. Insulating layer

[0042] An insulating layer is applied to the active layer in order to separate the layer containing the benzimidazole(s) from the enteric coating layer. In fact, gastroresistant polymers generally have free carboxyl groups with acidic characteristics, meaning that if they were to come into direct contact with the benzimidazoles they would cause the degradation thereof. For this reason, an insulating layer should be inserted between the active layer and the enteric coating layer.

[0043] The insulating layer contains binding agents of a polymeric nature which, by virtue of the evaporation of the solvent (i.e. demineralized water), form a protective layer. The binding agents of the insulating layer should be chosen among hydroxypropylmethylcellulose (Pharmacoat® of Shinetsu; Methocel® of Colorcon), hydroxypropylcellulose (Klucel® of Hercules), hydroxymethylcellulose, hydroxyethylcellulose (Natrosol® of Hercules or Cellosize® of Union Carbide Corporation), methylcellulose (Methocel® A of Colorcon), ethylcellulose (Aquacoat® de FMC or Surelease® of Colorcon), polyvinylpyrrolidone (Povidone® of GAF Corp., Kollidon® of BASF), polyethylene glycols, etc. The dispersion obtained, of suitable viscosity, is applied to the granules already containing the active ingredient layer. The thickness of this layer will not be less than 15 μm . Experimentally, it was ascertained that when this layer is not thick enough to suitably protect the layer containing the active ingredient(s) from contact with the enteric coating layer, the product is degraded quickly and darkens, and the gastroresistance of the final product quickly reaches values lower

than 85%. Thus, the nature and thickness of this intermediate layer was found to be decisive to the stability of the final product. The composition of this layer, of a strictly polymeric nature, provided that it is of appropriate thickness, prevents the incorporation of alkaline and alkaline-earthly metallic salts into the intermediate layer in order to ensure the stability of the product.

IV. Gastroresistant (or enteric) coating layer

[0044] In the pharmaceutical dosage form, the enteric coating layer is applied to the insulating layer and it contains anionic copolymers of methacrylic acid and ethyl acrylate (Eudragit® L30D, Eudragit® L30D-55, Eudragit® L100-55 of Rohm & Hass), or latex of cellulose acetophtalate (CAP), such as Aquateric® of FMC, pharmaceutically acceptable plasticizers, such as triethylcitrate (Citroflex®-2), tributylcitrate (Citroflex®-4), acetyltributylcitrate (Citroflex®-A4), dibutyl sebacate (DBS), diethylphtalate (DEP), acetylated monoglyceride (Myvacet® 9-40), polyethylene glycols or 1,2-propylene glycol, and possibly a lubricant such as talc or colloidal silicon dioxide (Aerosil® 200 of Degussa) and an anti-foaming agent in the nature of silicone. The amounts of polymer and plasticizer have to be strictly selected in order to guarantee the intended gastroresistance.

[0045] The gastroresistant film coats each granule completely and makes the dosage form insoluble in acid media, but quickly disintegrable for neutral and alkaline pH values, as in the case of the fluids present in the proximal fraction of the small intestine, where the dissolution and the absorption of the benzimidazole(s) will occur. The thickness of this layer should be not be less than 30 µm, in order to guarantee effective protection.

[0046] The final product is sieved in rooms with controlled humidity and temperature, and placed in hard gelatine capsules.

[0047] The dosage forms of the present invention are multi-unitary formulations in the form of pellets characterized by: a) a spherical inert core coated with three concentric layers which results in a product of spherical symmetry, b) a first layer of a thickness of 150-250 µm, applied to the inert core, containing at least one benzimidazole or a pharmaceutically acceptable salt thereof, mixed with inert substances that are also pharmaceutically acceptable, c) a second layer (intermediate layer), of a strictly polymeric nature and a minimum thickness of 15 µm, and d) a third layer (the outside layer), constituted by a gastroresistant or enteric polymer, of a minimum thickness of 30 µm. Moreover, the formulations are characterized by the complete absence of residues of organic solvents and the impurities associated to them, a characteristic which, among others, distinguishes them from the other formulations on the market.

[0048] Below, there are some examples of formulations that were obtained during the optimization process until the final formulations were produced.

EXAMPLES:

[0049]

Table 1 -

Qualitative and quantitative composition of dosage forms containing omeprazole (90±10 mg of omeprazole /g of pellets) according to the invention.										
COMPOSITION	Example 1	Example 2	Example 3	Example 4	Example 5	Example 6	Example 7	Example 8	Example 9	Example 10
Inert cores	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0	(g) 1000.0
<u>Active layer:</u>										
Omeprazole	140	140	140.0	140.0	140.0	140.0	140.0	140.0	140.0	140.0
Sodium dodecyl sulfate	2.7	2.7	4.0	4.0	4.0	4.0	4.0			
Sodium starch glycolate	4.3	4.3	4.3	4.3	4.3	4.3	4.3	4.0	4.0	4.0
Hydroxypropyl methylcellulose	11.0	11.0	11.0	11.0	11.0	11.0	11.0	56.3	56.3	56.3
Sodium Tetraborate	29.5	29.5	29.5	29.5	29.5	29.5	29.5	1.5	1.8	1.8
Lactose	63.2	47.2	45.3	45.3	45.3	45.3	45.3			
Emulsion 30% Simethicone	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Glycerol polyethylene glycoloxystearate	-	-	-	-	-	-	-	8.3	8.3	8.3
Talc	-	-	-	-	-	-	-	-	2.5	-
<u>Insulating layer:</u>										
Hydroxypropyl methylcellulose	40	40.0	40.0	40.0	46.7	80.0	80.0	80.0	80.0	80.0
<u>Enteric layer:</u>										
Copolymer of methacrylic acid	150	225.0	250.0	258.7	555.0	225.0	150.0	225.0	225.0	225.0
Triethylcitrate	15.0	22.5	25.0	25.9	55.0	22.5	15.0	22.5	22.5	22.5
Emulsion 30% Simethicone	0.9	1.5	1.7	1.8	3.8	1.5	1.0	1.5	1.5	1.5

Table I - (continued)

Qualitative and quantitative composition of dosage forms containing omeprazole (90 ± 10 mg of omeprazole /g of pellets) according to the invention.										
COMPOSITION	Example 1	Example 2	Example 3	Example 4	Example 5	Example 6	Example 7	Example 8	Example 9	Example 10
Enteric layer:										
TOTAL	1457.5	1524.6	1551.7	1561.4	1895.5	1564.0	1481.0	1540.0	1542.8	1540.3
Assay mg/g (theoretical value)	96.1	91.8	90.2	89.7	73.9	89.5	94.5	90.9	90.7	90.9
Tests	Example 1	Example 2	Example 3	Example 4	Example 5	Example 6	Example 7	Example 8	Example 9	Example 10
Assay (mg/g) (%)	79.6 (82.8%)	80.0 (87.1%)	78.5 (87.0%)	77.8 (86.7%)	67.8 (91.8%)	81.2 (90.7%)	85.6 (90.6%)	89.3 (98.2%)	85.5 (94.5%)	87.4 (96.1%)
Gastroresistance (%)	14.9	19.4	39.5	62.8	95.0	96.2	91.3	98.0	96.7	96.5
Dissolution (%)	-	-	-	-	84.3	93.3	86.3	91.8	96.5	96.3

[0050] The formulations presented in Examples 1 - 10 were obtained from inert cores coated successively by the three layers, each one with the composition presented in Table I.

[0051] The pharmaceutical formulations of the present invention take the form of pellets with which hard gelatine capsules are filled. The preparation of the pellets is processed as follows. For the preparation of the layer containing the active substance(s) (1st layer or active layer), a portion of demineralized water was heated up to 85-90°C and then hydroxypropylmethylcellulose (HPMC) was dispersed in it. In another portion of demineralized water, glycerol polypropyleneglycol oxystearate, sodium tetraborate, the simethicone emulsion, sodium starch glycolate and, finally, benzimidazole(s) were dispersed. When the suspension became uniform it was mixed with the HPMC dispersion after cooling to about 35°C. This last suspension was homogenized by shaking and applied to the inert cores (710 - 850 µm or 850 - 1000 µm). After the application of the suspension, the pellets were dried so that their temperature did not exceed 50°C.

[0052] For the preparation of the insulating layer (2nd layer or intermediate layer), demineralized water was heated up to 85-90°C. After this temperature was reached, HPMC was added and homogenized by shaking, until total dispersion was achieved. The dispersion was cooled to room temperature and applied to the product already containing the benzimidazolic layer. This operation is extremely delicate, because it must be guaranteed that the fluidization of the bed ensures the homogenous coating of all the pellets, with a film of regular thickness (which should not be less than 15 µm).

[0053] The gastroresistant or enteric coating layer (3rd layer or external layer) was prepared by mixing the demineralized water with the simethicone emulsion, followed by the addition of triethylcitrate and, finally, the copolymer of methacrylic acid. The dispersion was then homogenized for a few minutes and applied to the pellets already coated with the intermediate layer.

[0054] All the stages of the manufacture of the dosage forms took place in a fluid bed equipment with an inner partition device (wurster), under specific conditions.

[0055] As regards the final products, the amount of the active ingredient (mg of omeprazole / g of pellets), as well as the gastroresistance (%) and dissolution parameters of the formulations, were determined. The objective established for this invention was the obtention of dosage forms with the following characteristics: amount of active ingredient ≥ 85 mg / g of pellets, gastroresistance ≥ 95% and dissolution ≥ 85%, in order to establish the following specification limits (which should be maintained throughout the period of validity of the product): amount of active ingredient ≥ 80.75 mg / g of pellets, gastroresistance ≥ 85%, dissolution ≥ 75%.

[0056] The analysis of the results obtained in the assays performed with the different dosage forms in the Examples presented in Table I led to the following conclusions:

- a) The dosage form illustrated in Example 1 did not present enough gastroresistance.
- b) The dosage form illustrated in Example 2 is similar to that of Example 1, but the amount of enteric polymer was increased (3rd layer). This change was not enough to significantly improve the gastroresistance of the dosage form.
- c) Examples 3 and 4 illustrate dosage forms wherein, preponderantly, the amount of gastroresistant polymer was increased, but their gastroresistance do not reached the intended levels.
- d) Example 5 illustrates a formulation wherein the amount of enteric polymer was duplicated. At the same time, the thickness of the intermediate layer was increased slightly. Using this strategy it was possible to obtain pellets with the intended gastroresistance. However, the thickness of the gastroresistant layer seems to have a negative influence on the dissolution of the formulation.
- e) Example 6 illustrates a formulation wherein the amount of hydroxypropylmethylcellulose used in the 2nd layer was doubled and the amount of gastroresistant polymer used was the same as in Example 2. Under these conditions, a gastroresistance value of 96.2% and a dissolution value of 93.3% were obtained.
- f) Example 7 illustrates a dosage form wherein it was investigated whether maintaining the thickness of the 2nd layer made it possible to further reduce the amount of gastroresistant polymer to be applied. The results obtained in the gastroresistance and dissolution assays show that it is not advisable to reduce the amount of gastroresistant polymer beyond the values used in Examples 2 and 6.
- g) Examples 8, 9 and 10 illustrate dosage forms wherein an attempt was made to improve the yield of application of the 1st layer, without affecting the dissolution of the dosage form. In Example 8, sodium dodecyl sulphate was substituted by glycerol polyethyleneglycol oxystearate, the use of lactose was excluded and the percentage of hydroxypropylmethylcellulose (HPMC) was increased considerably. In Example 9, the amount of sodium tetraborate was corrected to adjust the pH value to about 9 and some talc was also included. Example 10 is similar to

Example 9, except that talc was not included as a lubricant.

h) The results obtained with the dosage forms illustrated in Examples 8, 9 and 10 suggest that the changes made to the composition of the 1st layer made it possible to increase by about 5% the yield of the industrial process for the application of the first layer, without affecting the dissolution of the dosage form.

[0057] As can be observed, Examples 1-5 and 7 are outside the scope of the present invention.

[0058] The Examples presented above showed that it is possible, using only aqueous solvents, to obtain formulations containing omeprazole which obey the dissolution and gastroresistance criteria required for these types of dosage forms. Examples 1 - 5 and 7 describe dosage forms which are executable from a technological point of view, but which do not obey the other criteria required for the type of dosage form intended to be developed. Examples 6, 8 - 10 describe dosage forms that obey the established criteria (i.e. GR > 95% and dissolution \geq 85%, after production); however, the dosage forms illustrated in examples 5 - 7 are economically not very advantageous, because the yield for application of the 1st layer, the one that contains omeprazole was still relatively low (i.e. about 90%), taking into account the high price of this active ingredient. Examples 8 - 10 constitute steps for optimizing the dosage forms in order to improve the yield of the industrial process, as a way of making it economically more advantageous and maintaining the intended characteristics of release and stability.

[0059] The present invention also relates to the preparation of other omeprazole multi-unitary dosage forms, of greater potency (i.e. 140 ± 10 mg of omeprazole/g of pellets), for treatment with higher dosages (i.e. 40 mg/capsule), also using hard gelatine capsules of small dimensions for increased patient comfort. In order to achieve this, inert cores of smaller dimensions are used (600-710 μ m). Two Examples of these formulations are presented as follows:

Table II -

Qualitative and quantitative composition of dosage forms containing omeprazole (140 ± 10 mg of omeprazole/g of pellets) according to the invention.		
COMPOSITION	Example 11	Example 12
	(g)	(g)
Inert cores	1000.0	1000.0
Active layer:		
Omeprazole	280.0	280.0
Sodium starch glycolate	8.0	8.0
Hydroxypropylmethylcellulose	112.6	112.6
Sodium tetraborate	3.0	3.0
Emulsion 30% Simethicone	1.8	1.8
Glycerol polyethyleneglycol oxystearate	16.3	16.3
Talc	-	5.0
Insulating layer:		
Hydroxypropylmethylcellulose	140.0	140.0
Enteric layer:		
Copolymer of methacrylic acid	390.0	390.0
Triethylcitrate	39.0	39.0
Emulsion 30% Simethicone	2.6	2.6
TOTAL	1993.3	1998.3
Assay mg/g (theoretical value)	140.5	140.1
Tests	Example 11	Example 12
Assay (mg/g) (%)	133.6 (95.1%)	134.8 (96.2%)

Table II - (continued)

Qualitative and quantitative composition of dosage forms containing omeprazole (140±10 mg of omeprazole/g of pellets) according to the invention.		
Tests	Example 11	Example 12
Gastroresistance (%)	96.1	95.0
Dissolution (%)	89.7	85.4

[0060] The preparation of the suspensions and dispersions for the manufacture of the product was carried out in the same manner as described previously.

[0061] In the case of these higher potency omeprazole dosage forms, since the inert cores have smaller dimensions, it was necessary to adjust the amount of hydroxypropylmethylcellulose to be incorporated into the intermediate layer in order to guarantee adequate insulation (i.e. thickness greater than 15 µm). It is also necessary to study the amount of enteric polymer coating to include in the formulation in order to obtain a layer with a thickness greater than 30 µm.

[0062] Just as it was possible to establish stable dosage forms for *omeprazole* (as shown later on), the present invention also seeks to describe multi-unitary dosage forms containing *lansoprazole*. Some Examples are presented below.

[0063] The preparation of the suspensions and dispersions was carried out according to the methodology described previously.

[0064] All the formulations illustrated in the Examples present assay, gastroresistance and dissolution parameters compatible with the requirements. Products manufactured in accordance with these Examples were submitted to long-term stability assays, the results of which are presented below.

Table III -

Qualitative and quantitative composition of dosage forms containing lansoprazole (90±10 mg of lansoprazole /g of pellets) according to the invention.			
COMPOSITION	Example 13	Example 14	Example 15
	(g)	(g)	(g)
Inert cores	1000.0	1000.0	1000.0
Active layer:			
Lansoprazole	140	140	140.0
Glycerol polyethyleneglycol ox. ystearate	8.3	8.3	8.3
Sodium starch glycolate	4.0	4.0	4.0
Hydroxypropylmethylcellulose	56.3	56.3	56.3
Sodium tetraborate	1.5	1.5	1.8
Emulsion 30% Simethicone	0.9	1.3	2.0
Talc	-	-	-
Insulating layer:			
Hydroxypropylmethylcellulose	80.0	80.0	80
Gastroresistant layer:			
Copolymer of methacrylic acid	225.0	225.0	225.0
Triethylcitrate	22.5	22.5	22.5
Emulsion 30% Simethicone	1.5	1.5	1.5
TOTAL	1540.0	1540.4	1541.4

Table III - (continued)

Qualitative and quantitative composition of dosage forms containing lansoprazole (90±10 mg of lansoprazole /g of pellets) according to the invention.			
COMPOSITION	Example 13	Example 14	Example 15
<u>Gastroresistant layer:</u>			
Assay mg/g (theoretical value)	90.9	90.9	90.8
Tests	Example 13	Example 14	Example 15
Assay (mg/g) (%)	90.5 (99.6%)	89.0 (97.9%)	88.6 (97.6%)
Gastroresistance (%)	92.3	94.0	95.1
Dissolution (%)	84.8	85.0	82.7

[0065] The present invention also applies to the active ingredient **pantoprazole**, with which multi-unitary dosage forms were also developed similar to the ones obtained for the other substituted benzimidazoles (omeprazole and lansoprazole). One Example of this application is presented below.

Table IV -

Qualitative and quantitative composition of dosage forms containing pantoprazole (90±10 mg of pantoprazole /g of pellets) according to the invention.	
COMPOSITION	Example 16 (g)
Inert cores	1000.0
<u>Active layer:</u>	
Pantoprazole	140
Glycerol polyethylenoglycol oxystearate	8.3
Sodium starch glycolate	4.0
Hydroxypropyl methylcellulose	56.3
Sodium tetraborate	2.0
Emulsion 30% Simethicone	1.8
<u>Insulating layer:</u>	
Hydroxypropylmethylcellulose	80.0
<u>Gastroresistant layer:</u>	
Copolymer of methacrylic acid	225.0
Triethylcitrate	22.5
Emulsion 30% Simethicone	1.5
TOTAL	1541.4
Assay mg/g (theoretical value)	90.8
Tests	Example 16
Assay (mg/g) (%)	90.0 (99.1%)

Table IV - (continued)

Qualitative and quantitative composition of dosage forms containing pantoprazole (90±10 mg of pantoprazole /g of pellets) according to the invention.	
Tests	Example 16
Gastroresistance (%)	93.8
Dissolution (%)	87.6

[0066] The present invention can extend its application to other substituted benzimidazoles with therapeutic interest, for example leminoprazole and/or pariprazole. It is also applicable to substituted benzimidazole salts, benzimidazole associations or pharmaceutically acceptable salts thereof.

[0067] With optimized dosage forms - final dosage forms - it will be shown below that the products obtained, as well as obeying analytic specifications when analyzed immediately after manufacture, maintain their stability during a period of time appropriate for sale on the pharmaceutical market for therapeutic purposes.

CONTROL OF THE FINISHED PRODUCT

[0068] Parameters analyzed:

- *Assay*
- *Gastroresistance*
- *Dissolution*
- *Total impurities.*

[0069] The above mentioned parameters were determined by HPLC (High Performance Liquid Chromatography), with an RP-8 column of 150 mm x 46 mm, filled with particles of 5 µm, and spectrophotometric detection at $\lambda = 280$ nm. In order to determine the total impurities, an RP-8 column of 250 mm x 46 mm was used.

[0070] The gastroresistance assays were performed by submitting the dosage forms, for 2 hours, to the action of an acidic incubation medium of pH ≈ 1.2, at 37°C ± 0.5°C, with constant mixing at 100 r.p.m. ± 4 r.p.m., using the dissolution equipment described by USP.

[0071] The dissolution assays were performed by submitting the dosage forms, for 2 hours, to the action of an acidic incubation medium of pH ≈ 1.2, at 37°C ± 0.5°C, with constant mixing at 100 r.p.m. ± 4 r.p.m., followed by the addition of a dissolution incubation medium, maintaining the same conditions for 30 min, using the dissolution equipment described by USP.

STABILITY TESTS

[0072] To test the stability of the pharmaceutical dosage forms, stability tests were performed on batches of capsules containing the pellets manufactured according to the present invention. The pellets were placed inside hard gelatine capsules which were stored in blisters of polyamide/aluminium/PVC sealed with aluminium foil of 25 µm. These blisters are maintained at room temperature (real time), and also in an incubator at 40°C and 75% relative humidity, and certain parameters were periodically determined, such as assay, gastroresistance, dissolution and total impurities. The results obtained for one batch of omeprazole pellets and for one batch of lansoprazole pellets are presented as follows.

Table V-

BATCH MED59 - OMEPRAZOLE PELLETS							
Assays performed with product maintained at 40°C/75% relative humidity							
Incubation period (months)	Temperat. (°C)	Relative Humidity (%)	Colour	Assay (mg/g) (%)	Gastro-resistance (%)	Dissolution (%)	Impurities Total Individual (%)
0	40	75	A	88.9 (100.0)	96.3	93.6	< 2.0 < 0.5

10

15

A-White Pearl

* The World Health Organisation considers that accelerated stability studies using incubation conditions of a temperature of 40°C and 75% relative humidity for 3 months in temperate and subtropical climates and 6 months in hot climates for products that are intended for the global market are a good indicator of the stability of a pharmaceutical dosage form. (WHO/PHARM/94. 565).

30

35

A-White Pearl.

Table VII-

BATCH LAN1- LANSOPRAZOLE PELLETS							
Assays performed with product maintained at 40°C/75% relative humidity							
Incubation period (months)	Temperat. (°C)	Relative Humidity (%)	Colour	Assay (mg/g) (%)	Gastro-resistance (%)	Dissolution (%)	Impurities Total partial (%)
0	40	75	A	90.5 (100.0)	90.9	83.1	<2.0 <0.5
1	40	75	A	90.5 (100.0)	92.1	82.9	<2.0 <0.5
3	40	75	A	86.3 (95.4)	92.9	81.2	<2.0 <0.5
6	40	75	A	87.1 (96.2)	91.2	80.7	<2.0 <0.5
<i>A-White Pearl</i> * The World Health Organisation considers that accelerated stability studies using incubation conditions of a temperature of 40°C and 75% relative humidity for 3 months in temperate and subtropical climates and 6 months in hot climates for products that are intended for the global market are a good indicator of the stability of a pharmaceutical dosage form. (WHO/PHARM/94.565).							

Table VIII-

BATCH LAN1- LANSOPRAZOLE PELLETS							
Assays performed with product maintained at room temperature (real time)							
Incubation period (months)	Temperat. (°C)	Relative Humidity (%)	Colour	Assay (mg/g) (%)	Gastro-resistance (%)	Dissolution (%)	Impurities Total partial (%)
0	25	60	A	90.5 (100.0%)	90.9	83.1	<2.0 <0.5
6	25	60	A	90.5 (100.0%)	89.1	82.0	<2.0 <0.5
9	25	60	A	-	-	-	-
12	25	60	A	90.3 (99.8%)	92.3	84.8	<2.0 <0.5

DISCUSSION

[0073] The Examples presented above show that the dosage forms of the present invention present good stability during a long period of storage (i.e. at least 3 years) and present gastroresistance parameters suitable for the length of time that the product stays in the stomach, being quickly dissolved as soon as it reaches the proximal portion of the small intestine. For these reasons, the formulations should present the following characteristics:

1) An inert core of spherical symmetry constituted by pharmaceutically acceptable inert excipients, which serves as the support on which the remaining elements of the formulation are deposited.

2) A layer containing at least one benzimidazole (i.e. omeprazole, lansoprazole or pantoprazole or other substituted benzimidazoles of therapeutic interest), or their various pharmaceutically acceptable salts, mixed in an appropriate proportion with pharmaceutically acceptable inert excipients, whose properties guarantee the disaggregation of the formulation and the dissolution of the active ingredient in the chosen site.

3) An insulating layer of a strictly polymeric nature, water-soluble, free from alkaline or alkaline-earthly metallic salts, of a minimum thickness of 15 μm , which separates the active layer from the gastroresistant or enteric layer.

4) An enteric coating layer, generally constituted by an acrylic polymer, of a minimum thickness of 30 μm , possibly in the presence of a plasticizer and/or an anti-foaming agent, insoluble at the gastric pH and easily disaggregable at the pH of the proximal part of the small intestine.

5) Total absence of residues of organic solvents or the impurities associated to them, since the formulation is produced exclusively using demineralized water.

[0074] The results obtained show that the dosage forms of the present invention are stable in terms of dosage as well as in terms of the other tested parameters (i.e. gastroresistance, dissolution and impurities). The process used to manufacture the products is economically more advantageous than the other inventions which belong to the current state of the art, and it also presents novelty as regards its applications. The technological process used is incomparably safer than the ones already described, since it does not require the use of organic solvents, with a toxicity risk to operators or an explosion risk. Moreover, the pharmaceutical dosage form itself is safer, since there are no residues of organic solvents or the impurities inherent to them in the finished products, which is an advantage of these products in relation to the others in terms of public health.

[0075] In addition, the whole manufacturing process takes place in the same equipment - a fluid bed equipment with an inner partition device (wurster), which produces pellets of high sphericity and a flat surface, constituted by various layers of uniform thickness, contrary to what happens with the pellets produced by extrusion/spheronization, by ro-togranulation or in conventional coating pans.

[0076] The pharmacist can always make variations to the pellet formulations of the present invention, for example the use of other aqueous polymers to ensure gastroresistance, such as aqueous dispersions of cellulose acetophthalate (CAP), or other plasticizers such as tributylcitrate (Citroflex®-4), acetyltributylcitrate (Citroflex®-A4), dibutyl sebacate (DBS), diethylphthalate (DEP), acetylated monoglyceride (Myvacet® 9-40), of other aggregants such as, methylcellu-lose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, polyethylene glycol, polyvinylpyrrolidone, etc., of other disaggregants such as pregelatinized starch, microcrystalline cellulose, calcic or sodic carboxymethylcellulose or sodium alginate, or other surfactant agents such as sodium trilaurylphosphate, sodium tridecylethoxylate, sodium myristylsulfate or sorbitane esters without going beyond the spirit and scope of the present invention. The invention can also be applied to benzimidazolic associations or pharmaceutically acceptable salts thereof, if this can be justified in therapeutic terms.

[0077] Finally, it should be pointed out that the present invention does not include formulations obtained by the classic extrusion/spheronization techniques already known from the state of the art, the products of which, in terms of quali-tative and quantitative composition and also structurally, are different from the ones that the present invention intends to protect.

Claims

1. Stable pellet preparations suitable for filling hard gelatine capsules, containing substituted benzimidazole(s) or their salts, **characterized in that** they contain a quantity of active ingredient(s) of between 1 - 50 mg an inert core of spherical symmetry with a diameter of 600 - 1000 μm constituted by inert excipients, coated by three sequential concentric layers as follows:

a) an active layer of a thickness of 150 - 250 μm containing at least one substituted benzimidazole and various pharmaceutically acceptable inert excipients, mixed in suitable proportions in order to allow a quick disaggre-gation of the formulations and dissolution of the active ingredient(s) in the proximal fraction of the small intes-tine;

b) an insulating layer of strictly polymeric nature, soluble in water, free from alkaline and/or alkaline-earthly metallic salts, of a minimum thickness of 15 μm ;

c) a gastroresistant or enteric layer of a minimum thickness of 30 μm ;

obtainable by a process in which all the stages of their manufacture take place in a fluid bed dryer equipment with wurster insert, without using organic solvents in any of the phases of the technological process.

2. Pharmaceutical preparations according to claim 1, **characterized in that** the layer which contains the active ingredient(s) comprises the alkaline compound sodium tetraborate.
3. Pharmaceutical preparations according to claims 1 - 2, **characterized in that** the layer which contains the active ingredient(s) comprises disaggregant agent(s), such as sodium starch glycolate, pregelatinized starch, microcrystalline cellulose, sodic or calcic carboxymethylcellulose or sodium alginate.
4. Pharmaceutical preparations according to claims 1 - 3, **characterized in that** layer which contains the active ingredient(s) comprises the surfactant glycerol polyethyleneglycol oxystearate.
5. Pharmaceutical preparations according to claims 1 - 4, **characterized by** an insulating layer which is soluble in water, free from of alkaline and/or alkaline-earthly metallic salts, containing exclusively binder agent(s) of polymeric nature in a percentage between 5 - 10% in weight (in relation to the final weight of the product), chosen among hydroxymethylcellulose, hydroxypropylcellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone or polyethylene glycols, in the absence of any compound other than residual water.
6. Pharmaceutical preparations according to claims 1-5, **characterized in that** the gastroresistant or enteric layer contains only aqueous polymers, such as the anionic co-polymers of methacrylic acid and ethyl acrylate, or latex of cellulose acetophthalate, present in a percentage between 14 - 20% in weight, in relation to the final weight of the product.
7. Pharmaceutical preparations according to claim 6, **characterized in that** the gastroresistant or enteric layer contains one or more plasticizers acceptable from a pharmaceutical point of view, such as triethylcitrate or polyethylenoglycols, mixed with an anti-foaming agent in the nature of silicone.

Patentansprüche

1. Stabile Pellet-Präparate, die zum Füllen von Hartgelatine kapseln geeignet sind und substituierte(s) Benzimidazol (e) oder ihre Salze enthalten, **dadurch gekennzeichnet, dass** sie eine Wirkstoffmenge von 1 - 50 mg und einen inerten Kern enthalten, der von kugelförmiger Symmetrie ist, einen Durchmesser von 600 - 1000 µm hat, durch inerte Arzneistoffträger gebildet wird und wie folgt mit drei aufeinander folgenden konzentrischen Schichten beschichtet ist:

- (a) eine aktive Schicht mit einer Dicke von 150 - 250 µm, die mindestens ein substituiertes Benzimidazol und verschiedene pharmazeutisch akzeptable inerte Arzneistoffträger enthält, gemischt in Verhältnissen, die geeignet sind, eine schnelle Aufschließung der Formulierungen und Auflösung des Wirkstoffs (der Wirkstoffe) im oberen Teil des Dünndarms zu erlauben;
- (b) eine isolierende Schicht von rein polymerer Natur, in Wasser löslich, frei von Alkali- und/oder Erdalkalimetallsalzen, mit einer Mindestdicke von 15 µm;
- (c) eine magensaftresistente oder enterische Schicht mit einer Mindestdicke von 30 µm;

erhältlich durch ein Verfahren, in dem alle Stadien ihrer Herstellung in einer Wirbelschichttrockner-Anlage mit Wurster-Einsatz ohne Verwendung von organischen Lösungsmitteln in irgendeiner der Phasen des technischen Verfahrens stattfinden.

2. Pharmazeutische Präparate nach Anspruch 1, **dadurch gekennzeichnet, dass** die Schicht, die den (die) Wirkstoff (e) enthält, die alkalische Verbindung Natriumtetraborat umfasst.
3. Pharmazeutische Präparate nach den Ansprüchen 1 - 2, **dadurch gekennzeichnet, dass** die Schicht, die den (die) Wirkstoff(e) enthält, Tablettensprengmittel wie zum Beispiel Natriumstärkeglycolat, vorgelatinisierte Stärke, mikrokristalline Cellulose, Natrium- oder Calciumcarboxymethylcellulose oder Natriumalginat, umfasst.
4. Pharmazeutische Präparate nach den Ansprüchen 1 - 3, **dadurch gekennzeichnet, dass** die Schicht, die den (die) Wirkstoff(e) enthält, das Tensid Glycerinpolyethylenglycoloxystearat umfasst.
5. Pharmazeutische Präparate nach den Ansprüchen 1 - 4, **gekennzeichnet durch** eine isolierende Schicht, die in Wasser löslich und frei von Alkali- und/oder Erdalkalimetallsalzen ist und ausschließlich Bindemittel polymerer

Natur, ausgewählt aus Hydroxymethylcellulose, Hydroxypropylcellulose, Hydroxypropylmethylcellulose, Polyvinylpyrrolidon oder Polyethylenglycolen, bei Fehlen jeder anderen Verbindung außer Restwasser, in einem Prozentsatz von 5 - 10 Gewichtsprozent (bezogen auf das Endgewicht des Produkts) enthält.

6. Pharmazeutische Präparate nach den Ansprüchen 1 - 5, **dadurch gekennzeichnet, dass** die magensaftresistente oder enterische Schicht nur wässrige Polymere, wie zum Beispiel die anionischen Copolymere von Methacrylsäure und Ethylacrylat, oder Latex von Celluloseacetatphthalat enthält, die in einem Prozentsatz von 14 - 20 Gewichtsprozent bezogen auf das Endgewicht des Produkts vorliegen.
7. Pharmazeutische Präparate nach Anspruch 6, **dadurch gekennzeichnet, dass** die magensaftresistente oder enterische Schicht einen oder mehrere Weichmacher, die vom pharmazeutischen Standpunkt her akzeptabel sind, wie zum Beispiel Triethylcitrat oder Polyethylenglycole, vermischt mit einem Antischaummittel in der Art von Silicon, enthält.

Revendications

1. Préparations de pastilles stables destinées au remplissage de capsules en gélatine dure, contenant un ou des benzimidazoles substitués ou leurs sels, **caractérisées en ce qu'elles** contiennent une quantité de principe(s) actif(s) entre 1 - 50 mg, un noyau inerte de symétrie sphérique ayant un diamètre de 600 - 1000 μm , constitué par des excipients inertes, revêtu par trois couches concentriques séquentielles comme suit :

a) une couche active d'une épaisseur de 150 - 250 μm contenant au moins un benzimidazole substitué et différents excipients inertes, acceptables d'un point de vue pharmaceutique, mélangés en proportions appropriées pour permettre une désaggrégation rapide des formulations et la dissolution du ou des principes actifs dans la fraction proximale du petit intestin ;

b) une couche isolante de nature strictement polymère, soluble dans l'eau, exempte de sels métalliques alcalins et/ou alcalino-terreux, d'une épaisseur minimum de 15 μm ;

c) une couche gastrorésistante ou entérique d'une épaisseur minimum de 30 μm ;

pouvant être obtenues par un procédé dans lequel toutes les étapes de leur fabrication s'effectuent dans un appareil de séchage à lit fluide avec un insert wurster, sans utiliser de solvants organiques dans l'une quelconque des phases du procédé technologique.

2. Préparations pharmaceutiques selon la revendication 1, **caractérisées en ce que** la couche qui contient le ou les principes actifs comprend le composé alcalin tétraborate de sodium.
3. Préparations pharmaceutiques selon les revendications 1 - 2, **caractérisées en ce que** la couche qui contient le ou les principes actifs comprend un ou des agents désagrégeants, tels que du glycolate d'amidon de sodium, d'amidon pré - gélatinisé, de cellulose microcristalline, de carboxyméthylcellulose sodique ou calcique ou d'alginate de sodium.
4. Préparations pharmaceutiques selon les revendications 1 - 3, **caractérisées en ce que** la couche qui contient le ou les principes actifs comprend l'agent tensioactif oxystéarate de glycérol polyéthylène - glycol.
5. Préparations pharmaceutiques selon les revendications 1 - 4, **caractérisées par** une couche isolante qui est soluble dans l'eau, exempte de sels métalliques alcalins et/ou alcalino-terreux, contenant exclusivement un ou des agents liants de nature polymère dans un pourcentage entre 5 - 10 % en poids (par rapport au poids final du produit), choisis parmi l'hydroxyméthylcellulose, l'hydroxypropylcellulose, l'hydroxypropyl méthylcellulose, la polyvinylpyrrolidone ou les polyéthylène - glycols, en l'absence de tout composé autre que de l'eau résiduelle.
6. Préparations pharmaceutiques selon les revendications 1 - 5, **caractérisées en ce que** la couche gastrorésistante ou entérique ne contient que des polymères aqueux, tels que les copolymères anioniques de l'acide méthacrylique et d'éthylacrylate, ou du latex de cellulose acétophthalate, présents dans un pourcentage entre 14 - 20 % en poids, rapporté au poids final du produit.

7. Préparations pharmaceutiques selon la revendication 6, **caractérisées en ce que** la couche gastrorésistante ou entérique contient un ou plusieurs agents plastifiants acceptables du point de vue pharmaceutique, tels que du triéthylcitrate ou des polyéthylèneglycols, mélangés avec un agent anti-moussant de la nature de la silicone.

5

10

15

20

25

30

35

40

45

50

55

Graphic representation of the results obtained for the stability tests carried out at 40°C/75% relative humidity with a batch of omeprazole pellets.




Figure 1

Graphic representation of the results obtained for the stability tests carried out in real time with a batch of omeprazole pellets.




Figure 2

Graphic representation of the results obtained for the stability tests carried out at 40°C/75% relative humidity with a batch of lansoprazole pellets.




Figure 3

Graphic representation of the results obtained for the stability tests carried out in real time with a batch of lansoprazole pellets.




Figure 4